



200278-04

**For the Patient**

***Cordran<sup>®</sup> Tape***  
***Flurandrenolide Tape, USP***

**DIRECTIONS FOR USE**

**APPLICATION OF MEDICATED TRANSPARENT TAPE**

Rx only


Revised: January 2008

**IMPORTANT:** Skin should be clean and *dry* before tape is applied. Tape should always be cut, never torn.

1. Prepare skin as directed by your physician or as follows: Gently clean the area to be covered to remove scales, crusts, dried exudates, and any previously used ointments or creams. A germicidal soap or cleanser should be used to prevent the development of odor under the tape. Shave or clip the hair in the treatment area to allow both good contact with the skin and comfortable removal. If a shower or tub bath is to be taken, it should be completed before the tape is applied. The skin should be dry before application of the tape.
2. Remove tape from package and cut a piece slightly larger than area to be covered. Round off corners.

**Cordran<sup>®</sup> Tape**  
*Flurandrenolide Tape, USP*

3. Pull white paper backing from transparent tape. Be careful that tape does not stick to itself.



4. Apply tape, keeping skin smooth; press tape into place.




**REPLACEMENT OF TAPE**

Unless instructed otherwise by your physician, replace tape after 12 hours. Cleanse skin and allow it to dry for 1 hour before applying new tape.

**DISPENSING THE TAPE**

To use the roll as a dispenser, pull the tape as illustrated.



**IF IRRITATION OR INFECTION DEVELOPS,  
REMOVE TAPE AND CONSULT PHYSICIAN.**

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F).

**Keep out of reach  
of children.**

**Rx only**

Revised: January 2008

S0108

**Watson<sup>®</sup>** 

Manufactured for:  
Watson Pharmaceuticals, Inc.  
Corona, CA 92880 USA

Manufactured by:  
3M Company  
St. Paul, MN 55144 USA